

BIOCOMPATIBLE PHOTOACOUSTIC NANOPARTICULAR CONTRAST AGENTS BASED ON BODIPY-SCAFFOLD AND POLYLACTIDE POLYMERS

Jean-Baptiste Bodin^{1,3}, Justine Cois², Flora Lefebvre³, Magali Noiray³, Gilles Clavier², Jérôme Gateau⁴, Nicolas Tsapis³, Rachel Méallet-Renault^{a1}

¹Institut des Sciences Moléculaires d'Orsay (ISMO), CNRS, Université Paris-Saclay, 91405 Orsay, France

² Laboratoire de Photophysique et Photochimie Supramoléculaires et Macromoléculaires (PPSM), Université Paris-Saclay, ENS Paris-Saclay, CNRS, , 91190, Gif-sur-Yvette, France

³ Institut Galien Paris-Saclay, Université Paris-Saclay, CNRS, 92296, Châtenay-Malabry, France

⁴ Institut Langevin, ESPCI, CNRS UMR 7587, 1 rue Jussieu, 75238 Paris cedex 05, France

e-mail address : jean-baptiste.bodin@universite-paris-saclay.fr

Photoacoustic imaging is an emerging biomedical imaging modality combining optical and ultrasound waves to map optical-absorption contrast at centimetric depth with sub-millimeter resolution. The key is the photoacoustic (PA) effect: optically absorbing structures emit ultrasound waves when excited with a ns-laser pulse. To reach cm-depth, PA imaging operates in the near-infrared (NIR) window in biological tissue (650-1000nm). NIR optical absorbers can thus be mapped throughout the range of depths and resolution explorable with medical ultrasound. We have designed novel PA molecules based on the BODIPY scaffold. These PA-BODIPYs were used as initiators for the ring opening polymerization of lactide to yield BODIPY-polylactide, that were further formulated into nanoparticles (NP). We present here the full spectroscopic and photoacoustic characterizations of the PA-BODIPYs, the corresponding polymers and NPs. Results show BODIPY NPs are promising contrast agents for PA imaging.




Figure 1 : From Bodipy to nanoparticles.

References:

- [1] J. Weber, P. C. Beard, S. E. Bohndiek. *Nature Methods*, 2016, 13 (8), 639-50.
- [2] Borg RE, Rochford J. *Photochem Photobiol.*, 2018 Nov, 94(6),1175-1209.
- [3] Y Ni, RK Kannadorai, SWK Yu, YT Chang, *Org. Biomol. Chem.*, 2017, 15, 4531-4535.