

EXTENDED LABORATORY INVESTIGATION OF THE PURE ROTATIONAL SPECTRUM OF THE CH₂CN RADICAL IN THE (SUB-)MILLIMETER REGION (79-860 GHz)

OLIVIA CHITARRA, THOMAS SANDOW HEARNE, OLIVIER PIRALI, MARIE-ALINE MARTIN-DRUMEL, *Institut des Sciences Moléculaires d'Orsay, Université Paris Saclay, CNRS, Orsay, France.*

The cyanomethyl radical, CH₂CN, is considered a key reactive intermediate in the interstellar medium (ISM) since its first detection [1]. To date, the radical has been detected in several environments of the ISM using pure rotational data available in the literature, limited to frequencies below 280 GHz [2,3]. The radical is also postulated to participate to the formation of complex organic molecules, such as cyanoacetaldehyde [4]. To enable the detection of the CH₂CN radical in current high frequency astronomical surveys, laboratory re-investigation of its spectrum at submillimeter wavelengths appears essential. We have investigated the pure rotational spectrum of CH₂CN at room temperature in the 75-900 GHz domain. The radical was produced using a H-abstraction method from CH₃CN using F atoms. To record pure rotational transitions, we used two spectrometers: a commercial broadband chirped-pulse (CP) spectrometer covering the 75-110 GHz spectral region and a tunable single-frequency absorption spectrometer exploiting a frequency multiplication chain with a large spectral coverage (here, 140-900 GHz). A combined fit of the literature data and our newly measured transitions (involving N'' and K_a'' up to 42 and 8, respectively) yields to an improvement of the rotational parameters; in particular the A rotational constant and K -dependent parameters. This work allows for confident searches of the radical in cold to warm environments of the ISM, over a wide frequency range. In addition, the broadband capacities of the CP spectrometer has also revealed very efficient in the study of discharge products (synthesized by the reaction between CH₃CN and F atoms in this work). I will present both aspects of this work: the improvement of the spectroscopy of CH₂CN and the analysis of the chemical composition of the discharge by CP spectroscopy.

[1] W. M. Irvine et al., *The Astrophysical Journal Letters* (1988) 334, L107 [2] S. Saito et al., *The Journal of Chemical Physics* (1997) 107, 1732 [3] H. Ozeki et al., *American Astronomical Society* (2004) 617, 680 [4] B. Ballotta et al., *ACS Earth and Space Chemistry* (2021) 5, 1071

NOTE: The horizontal line across the page indicates the maximum allowed length of the abstract including footnotes. If your abstract is close to or exceeds the length limit, it may appear on page 2 of this document.

Time required: 15 min

Session Categories (Keywords) by Relevance: Rotational structure/frequencies — Radicals — Chirped pulse — Astronomy

Mini-Symposia Requested: None — None

Competing for Rao Prize? Yes

Competing for Miller Prize? No